



Figure 1: Geometry used to define the volume scattering function. $\Phi_i(\lambda)$ is the incident power onto an area ΔA . An amount $\Delta\Phi_a(\lambda)$ of the incident power is absorbed in distance Δr , and $\Delta^2\Phi_s(\psi, \lambda)$ is the power scattered into the annular ring of solid angle $\Delta\Omega$. $\Delta\Phi_t(\lambda)$ is the transmitted power.

We now take into account the angular distribution of the scattered power, with two assumptions. We first assume that the medium is isotropic, i.e., its influence on light is the same in all directions at a given point. This is a reasonable assumption for natural waters in which the particles are randomly oriented by turbulence. We also assume that the incident light is unpolarized. If these two assumptions are true, then the scattering process is azimuthally symmetric. This means that the scattering depends only on the scattering angle ψ , which is measured from the direction of the unscattered beam as shown in Fig. figure1. Clearly $0 \leq \psi \leq \pi$.

Conceptual Definition of the Volume Scattering Function

With these two assumptions, the *fraction* of incident power scattered out of the beam through an angle ψ into a solid angle $\Delta\Omega$ centered on ψ , is $\Delta^2\Phi_s(\psi, \lambda)/\Phi_i(\lambda)$. The notation $\Delta^2\Phi_s(\psi, \lambda)$ is adopted from ? to indicate that $\Delta^2\Phi_s(\psi, \lambda)$ is a second-order small quantity, that is, it is small because both Δr and $\Delta\Omega$ are small. The solid angle $\Delta\Omega$ now includes all directions within the two red rings shown in the figure, corresponding to all directions between scattering angles ψ and $\psi + \Delta\psi$. The volume scattering function $\beta(\psi, \lambda)$ is defined as the limit of this fraction as $\Delta r \rightarrow 0$ and $\Delta\Omega \rightarrow 0$:

$$\beta(\psi, \lambda) \equiv \lim_{\Delta r \rightarrow 0} \lim_{\Delta\Omega \rightarrow 0} \frac{\Delta^2\Phi_s(\psi, \lambda)}{\Phi_i(\lambda) \Delta r \Delta\Omega} = \frac{1}{\Phi_i(\lambda)} \frac{\partial^2\Phi_s(\psi, \lambda)}{\partial r \partial\Omega} \quad (\text{m}^{-1} \text{sr}^{-1}). \quad (1)$$

Recalling the definition of *spectral radiant intensity* ?? as scattered power per unit solid angle, the corresponding intensity scattered into the given solid angle $\Delta\Omega$ is $\Delta I_s(\psi, \lambda) = \Delta^2\Phi_s(\psi, \lambda)/\Delta\Omega$. Moreover, if the incident power $\Phi_i(\lambda)$ falls on an area ΔA , then the corresponding incident irradiance is $E_i(\lambda) = \Phi_i(\lambda)/\Delta A$. Noting that $\Delta V = \Delta r \Delta A$ is the volume

of water that is illuminated by the incident beam allows Eq. (likesection1) to be rewritten as

$$\beta(\psi, \lambda) = \lim_{\Delta V \rightarrow 0} \frac{\Delta I_s(\psi, \lambda)}{E_i(\lambda) \Delta V} = \frac{1}{E_i(\lambda)} \frac{\partial I_s(\psi, \lambda)}{\partial V} . \quad (2)$$

This form of $\beta(\psi, \lambda)$ suggests the name *volume scattering function* (commonly abbreviated as VSF) and the physical interpretation of scattered intensity per unit incident irradiance per unit volume of water. In the language of a physicist, the VSF also can be interpreted as the differential scattering cross section per unit volume.

The forms of Eqs. (likesection1) and (likesection2) involving partial derivatives are *conceptual* definitions. Instruments that measure the VSF always have to deal with finite path lengths Δr and finite solid angles $\Delta\Omega$. The transition from a definition as a derivative to an instrument design is discussed on Sec. ?? on measuring IOPs.

Scattering Coefficients

Integrating $\beta(\psi, \lambda)$ over all directions (solid angles) gives the total scattered power per unit incident irradiance and unit volume of water, in other words the scattering coefficient:

$$b(\lambda) = \int_{\Xi} \beta(\psi, \lambda) d\Omega = 2\pi \int_0^\pi \beta(\psi, \lambda) \sin \psi d\psi . \quad (3)$$

The 2π in the last equation follows from our assumption that the scattering is azimuthally symmetric about the incident direction. This integration is often divided into forward scattering, $0 \leq \psi < \pi/2$, and backward scattering, $\pi/2 \leq \psi \leq \pi$, parts. The corresponding forward and backward scattering coefficients are, respectively,

$$b_f(\lambda) \equiv 2\pi \int_0^{\pi/2} \beta(\psi, \lambda) \sin \psi d\psi \quad (4)$$

$$b_b(\lambda) \equiv 2\pi \int_{\pi/2}^\pi \beta(\psi, \lambda) \sin \psi d\psi . \quad (5)$$

The backscatter fraction, defined by

$$B_b(\lambda) = \tilde{b}_b \equiv \frac{b_b(\lambda)}{b(\lambda)} , \quad (6)$$

gives the fraction of scattered light that is deflected through scattering angles greater than 90 deg. This quantity is fundamental to remote sensing because most of the light leaving the ocean in an upward direction comes from sunlight that was originally heading downward, but which was backscattered into upward directions.

The Scattering Phase Function

The volume scattering *phase function*¹, $\tilde{\beta}(\psi, \lambda)$, is defined by

$$\tilde{\beta}(\psi, \lambda) \equiv \frac{\beta(\psi, \lambda)}{b(\lambda)} \quad (\text{sr}^{-1}). \quad (7)$$

Writing the VSF $\beta(\psi, \lambda)$ as the product of the scattering coefficient $b(\lambda)$ times the phase function $\tilde{\beta}(\psi, \lambda)$ partitions $\beta(\psi, \lambda)$ into a factor giving the *magnitude* of the total scattering, $b(\lambda)$ with units of m^{-1} , and a factor giving the *angular distribution* of the scattered light, $\tilde{\beta}(\psi, \lambda)$, with units of sr^{-1} . Combining Eqs. (likesection3) and (likesection7) gives the *normalization condition for the phase function*:

$$2\pi \int_0^\pi \tilde{\beta}(\psi, \lambda) \sin \psi \, d\psi = 1. \quad (8)$$

This normalization implies that the backscatter fraction can be computed from

$$B_b(\lambda) \equiv 2\pi \int_{\pi/2}^\pi \tilde{\beta}(\psi, \lambda) \sin \psi \, d\psi. \quad (9)$$

The *asymmetry parameter* g or *mean cosine* of the phase function is average over all scattering directions of the cosine of the scattering angle ψ , namely

$$g = \langle \cos \psi \rangle \equiv 2\pi \int_0^\pi \tilde{\beta}(\psi) \cos \psi \sin \psi \, d\psi. \quad (10)$$

The asymmetry parameter is a convenient measure of the “shape” of the phase function. For example, if $\tilde{\beta}(\psi)$ is very large for small ψ , then g is near one. If $\tilde{\beta}(\psi)$ is symmetric about $\psi = 90$ degrees, then $g = 0$. Typical ocean waters have g values in the range of 0.8 to 0.95.

Comment: We have assumed above that the scattering is azimuthally symmetric, so that the directional pattern or angular shape of the VSF depends only on the scattering angle ψ . This is not the case for polarized incident light, even if the medium is isotropic. Thus a linearly polarized laser beam will scatter differently for different azimuthal directions (measured relative to the plane of polarization), even if the medium is isotropic. Description of polarized light requires the Stokes vector formalism and a more complicated description of scattering by polarized light; also see The General Vector RTE.

If the medium contains non-spherical particles that are not randomly oriented, even unpolarized light will scatter differently for different azimuthal directions. This is sometimes

¹The term “phase function” has its historical origins in astronomy, where the phase angle is the angle between the directions of the light incident onto and reflected from an object. In the case of sunlight being reflected by the Moon and seen on Earth, this angle depends on the phase of the Moon—hence the name phase function for the function that gives the phase angle as a function of time. In our terminology, the lunar phase angle is π minus the scattering angle. The term phase function is rather misleading because the phase function $\tilde{\beta}(\psi, \lambda)$ defined here has nothing whatsoever to do with the phase of an electromagnetic wave or with its historical origin tied to the phase of the Moon. The Russian literature calls $\tilde{\beta}$ the “scattering diagram,” which is more descriptive of what it really is. You also see the term “indicatrix.” Neither of these terms is commonly used in the English-language literature.

the case, for example, in cirrus clouds and ice fogs, which can have non-spherical ice crystals that become oriented in a particular way as the crystals fall through a calm atmosphere. The atmosphere is then an optically anisotropic medium, and scattering is not azimuthally symmetric. Scattering of sunlight in such an atmosphere gives rise to phenomena such as “sun dogs” or parhelia, which are bright spots to either side of the sun.